



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

Solution by G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va.

Let  $x=y+z$ . Then the equation becomes

$$y^7+z^7+(7yz+p)[y^5+z^5+(3yz+\frac{2}{7}p)(y^3+z^3) \\ + (5y^2z^2+\frac{5}{7}pyz+\frac{1}{43}p^2)(y+z)]+r=0.$$

Now  $x$  may be decomposed into two parts,  $y$  and  $z$ , in an infinite number of ways; we may, therefore, suppose  $y$  and  $z$  are such as to satisfy the condition  $7yz+p=0$ .

$$\therefore yz=-\frac{1}{7}p, y^7+z^7=-r.$$

$$\text{Let } y^7=a, z^7=b. \therefore a+b=-r, ab=-(\frac{1}{7}p)^7.$$

$$\therefore a \text{ and } b \text{ are the roots of the equation } u^2+ru-(\frac{1}{7}p)^7=0.$$

$$\therefore a=-\frac{1}{2}r+\sqrt{[\frac{1}{4}r^2+(\frac{1}{7}p)^7]}, b=-\frac{1}{2}r-\sqrt{[\frac{1}{4}r^2+(\frac{1}{7}p)^7]}.$$

Let  $\omega$  be an imaginary seventh root of unity, so that

$$\omega=\frac{1}{2}[A\pm\sqrt{(A^2-4)}], A=\frac{1}{3}\sqrt[3]{\frac{2}{3}}[\sqrt[3]{(1+3\sqrt{-3})}+\sqrt[3]{(1-3\sqrt{-3})}-\sqrt[3]{\frac{2}{3}}].$$

Then the required seven roots of the equation are

$$\sqrt[7]{a}+\sqrt[7]{b}, \quad \omega\sqrt[7]{a}+\omega^6\sqrt[7]{b}, \quad \omega^2\sqrt[7]{a}+\omega^5\sqrt[7]{b}, \quad \omega^3\sqrt[7]{a}+\omega^4\sqrt[7]{b}, \\ \omega^6\sqrt[7]{a}+\omega\sqrt[7]{b}, \quad \omega^5\sqrt[7]{a}+\omega^2\sqrt[7]{b}, \quad \omega^4\sqrt[7]{a}+\omega^3\sqrt[7]{b}.$$

188. Proposed by GUY SCHUYLER.

$$xy+ab=2ax, \quad x^2y^2+a^2b^2=2b^2y^2.$$

Solution by O. W. ANTHONY, Head of Mathematical Department, DeWitt Clinton High School, New York City.

By squaring the first equation and subtracting from the second we get  $y=xa/b$  and  $y=-2xa/b$ . Whence easily  $x=b$ ,  $\frac{1}{2}(-1\pm\sqrt{3})b$ .  $y=a(1\mp\sqrt{3})b/a$ .

Also solved by G. W. Greenwood, B. A., Professor of Mathematics and Astronomy, McKendree College, Lebanon, Ill.; G. W. Drake, Fayetteville, Ark.; Charles E. Barrett, Louisville, Ky.; H. F. MacNeish, A. B., Instructor in Mathematics in the University High School, Chicago, Ill.; L. E. Newcomb, Los Gatos, Cal.; G. B. M. Zerr, A. M., Ph. D., Parsons, W. Va., and J. Scheffer, Kee Mar College, Hagerstown, Md.

## GEOMETRY.

Problem 203 was also solved by Henry A. Converse, Ph. D., Instructor in Mathematics, Johns Hopkins University, Baltimore, Md.

208. Proposed by W. J. GREENSTREET, A. M., Editor of The Mathematical Gazette, Stroud, England.

Tangents drawn to two confocal parabolaes from the point on the common tangent intersect at the same angle as the axes of the parabolaes.

I. Solution by G. W. GREENWOOD, A. M. (Oxon), Professor of Mathematics and Astronomy in McKendree College, Lebanon, Ill.

$PQ$  is the common tangent;  $S$ , the common focus. Draw  $PA$ ,  $PA'$  parallel to the axes;  $PT$ ,  $PT'$  are the other tangents from  $P$ ;  $T$ ,  $T'$  being the points of tangency. Join  $PS$ .

Since a tangent from a point to a parabola makes the same angle with the line joining the point to the focus as the remaining tangent from the point does with the axis, we have  $\angle TPA = \angle QPS = \angle TPA'$ . Hence  $\angle TPT' = \angle APA'$ .

II. Solution by G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va.

$$\text{Let } y^2 = 4a(x+a) \dots (1),$$

$$y^2 \cos^2 \theta + 2y \sin \theta (x \cos \theta + 2b) = 4bx \cos \theta + 4b^2 - x^2 \sin^2 \theta \dots (2),$$

be the confocal parabola with axes inclined at an angle  $\theta$ .

$$\text{Let the common tangent be } y = mx + c \dots (3).$$

$$\text{If (3) is tangent to (1), } c = a(m^2 + 1)/m.$$

$$\text{If (3) is tangent to (2), } c = b(m^2 + 1)/(m \cos \theta + \sin \theta).$$

$$\therefore m = \frac{a \cos \theta}{b - a \cos \theta}, \quad c = \frac{a^2 + b^2 - 2ab \cos \theta}{\sin \theta (b - a \cos \theta)}.$$

$$\therefore (3) \text{ becomes } y = \frac{ax \sin \theta}{b - a \cos \theta} + \frac{a^2 + b^2 - 2ab \cos \theta}{\sin \theta (b - a \cos \theta)} \dots (4).$$

Let  $(h, k)$  be the point on (4).

$$\therefore y = m_1 x + a(m_1^2 + 1)/m_1 \text{ is tangent to (1),}$$

$$y = m_2 x + b(m_2^2 + 1)/(m_2 \cos \theta + \sin \theta) \text{ is tangent to (2).}$$

$$\text{But } k = m_1 h + a(m_1^2 + 1)/m_1 \dots (5),$$

$$k = m_2 h + b(m_2^2 + 1)/(m_2 \cos \theta + \sin \theta) \dots (6),$$

$$k = \frac{ah \sin \theta}{b - a \cos \theta} + \frac{a^2 + b^2 - 2ab \cos \theta}{\sin \theta (b - a \cos \theta)} \dots (7).$$

$$\text{From (5), } m_1^2 - \frac{km_1}{a+h} + \frac{a}{a+h} = 0 \dots (8).$$

$$\text{From (6), } m_2^2 + \frac{(h \sin \theta - k \cos \theta)m_2}{h \cos \theta + b} + \frac{b - k \sin \theta}{h \cos \theta + b} = 0 \dots (9).$$

$\therefore m_1$  and  $m_2$  each has two values.

From (4),  $m_1 = m_2 = a \sin \theta / (b - a \cos \theta)$ . The other values of  $m_1$  and  $m_2$  are respectively,

$$\frac{b - a \cos \theta}{(a + h) \sin \theta}, \quad \frac{(b - k \sin \theta)(b - a \cos \theta)}{a \sin \theta (h \cos \theta + b)} = \frac{b \cos \theta - h \sin^2 \theta - a}{\sin \theta (h \cos \theta + b)}.$$

$$\therefore \frac{m_1 - m_2}{1 + m_1 m_2} = \frac{(a^2 + b^2 - 2ab \cos \theta + 2ah \sin^2 \theta + h^2 \sin^2 \theta) \sin \theta}{(a^2 + b^2 - 2ab \cos \theta + 2ah \sin^2 \theta + h^2 \sin^2 \theta) \cos \theta} = \tan \theta.$$

209. Proposed by W. J. GREENSTREET, M. A., Editor of The Mathematical Gazette, Stroud, England.

Find by a geometrical method the maximum value of  $\sin \theta \cos \theta \cos 2\theta$ .